

Phosphorylation of Immunoglobulin-Containing and Proline-Rich Receptor-1 (IGPR-1): Identification and Functional Significance

Yan Jiang; Kobra Rezazadeh; Edward Hartsough; Rosana Meyer; Nader Rahimi; Catherine E. Costello

Boston University School of Medicine, Boston, MA

Background

IGPR-1 (Immunoglobulin containing and proline rich receptor (IGPR-1)) is a novel adhesion molecule expressed in endothelial cells and epithelial cells. IGPR-1 activity regulates cell-cell interaction and capillary tube formation/angiogenesis. The proline rich cytoplasmic domain of IGPR-1 strongly interacts with several SH3-containing signaling proteins including, SPIN90/WISH (SH3 protein interacting with Nck, 90 kDa/Wiskott–Aldrich syndrome protein (WASP) interacting SH3 protein). SPIN90 plays a central role in IGPR-1 mediated capillary tube formation of endothelial cells. To determine the molecular basis of interaction of IGPR-1 with SPIN90 and establish biological importance of phosphorylation to IGPR-1 function, we analyzed post-translational modifications of IGPR-1, particularly phosphorylation sites that might regulate its interaction with signaling proteins and its function.

Methods

Cells were prepared and lysed, and IGPR-1 was isolated by immunoprecipitation using anti-IGPR-1 antibody. On-bead reduction, alkylation and trypsin digestion were performed while immunoprecipitated proteins were still bound to the affinity beads. Enrichment of phosphopeptides was carried out using TiO₂. Separation on a nano C18 column was followed by MS/MS analysis (both CID and HCD) on an LTQ Orbitrap. LC-MS/MS data were analyzed using Proteome Discoverer. All the phosphopeptides assigned by Mascot and Sequest searches were verified manually.

Results and Conclusions

IGPR-1 is phosphorylated at three serine sites, including S220, S262 and S266. Site-directed mutagenesis analysis demonstrated that phosphorylation of S262 and S266 are not required for recognition of SPIN90 by IGPR-1, suggesting that perhaps phosphorylation of S220 but not S262, and S266, is important for interaction of IGPR-1 with SPIN90. Further studies are underway to elucidate the importance of S220 in the interaction of IGPR-1 with SPIN90 and the putative biological importance of S262 and S266 phosphorylations in IGPR-1 mediated cellular functions in milieu of angiogenesis.

This research is supported by NIH grants P41 RR010888/GM104603, S10 RR020946, R01 EY017955 and NIH-NHLBI contract N01 HV00239.