

Benzodiazepines

CRIT program
May 2011

Alex Walley, MD, MSc
Assistant Professor of Medicine
Boston University School of Medicine

Medical Director, Opioid Treatment Program
Boston Public Health Commission

Medical Director, Overdose Prevention Pilot Program
Massachusetts Department of Public Health

Learning objectives

At the end of this session, you should:

1. Understand why people use benzodiazepines
2. Know the characteristics of benzodiazepine intoxication and withdrawal syndromes
3. Understand the consequences of these drugs
4. Know the current options for treatment of benzodiazepine dependence

Roadmap

1. Case and controversies
2. History and Epidemiology
3. Benzo effects
4. Treatment



Case

- 29 yo man presents for follow-up at methadone clinic after inpatient admission for femur fracture from falling onto subway track from platform.
- Treated with clonazepam since age 16 for panic disorder. Takes 6mg in divided doses daily. Missed his afternoon dose, the day of the accident
- Started using heroin at age 23
- On methadone maintenance for 1 year, doing well, about to get his first take home

Thoughts

- Did BZDs cause his fall?
 - or did not taking BZDs cause his fall?
- Is he addicted to benzos?
- Should he come off of BZDs? How?
- If not, how do we keep him safe?
- Should he get take homes?
- Did teenage BZD treatment cause his heroin addiction?


History and Epidemiology



History

- First discovered in 1954 by a Austrian scientist, Leo Sternbach → Librium
- 1963 → Valium
- Used for anxiety, seizures, withdrawal, insomnia, drug-associated agitation

Valium® (diazepam)
effective at many levels of symptomatology



Psychic level
Valium (diazepam) helps reduce the conscious burden of tension that can interfere with work efficiency and enjoyment.

Psychosomatic level
Valium eases stress on the motor-visceral system, aiding medical management when somatic complaints are associated with emotional factors.

Somatic level
Valium, used adjunctively with other drugs or physiotherapy, favorably affects the entire cluster of skeletal muscle spasm-related symptoms.

Valium® (diazepam)
2-mg, 5-mg, 10-mg tablets to help relieve psychic tension and its somatic symptoms

Before prescribing, please consult complete product information, a summary of which follows.

Indications: Tension and anxiety states; somatic complaints which are concomitants of emotional factors; psychoneurotic states manifested by tension, anxiety, apprehension, fatigue, depressive symptoms or agitation; acute agitation, tremor, delirium, tremors and hallucinations due to acute alcohol withdrawal; adjunctively in skeletal muscle spasm due to reflex spasm to local pathology, spasticity caused by upper motor neuron disorders, athetosis, stiff-man syndrome, convulsive disorders (not for sole therapy).

Contraindicated: Known hypersensitivity to the drug. Children under 6 months of age. Acute narrow angle glaucoma.

Warnings: Not of value in psychotic patients. Caution against hazardous occupations requiring complete mental alertness. When used adjunctively in convulsive disorders, possibility of increase in frequency and/or severity of grand mal seizures may require increased dosage of standard anticonvulsant medication; abrupt withdrawal may be associated with temporary increase in frequency and/or severity of seizures. Advise against simultaneous ingestion of alcohol and other CNS depressants. Withdrawal symptoms have occurred following abrupt discontinuance. Keep addiction prone individuals under careful surveillance because of their predisposition to habituation and dependence. In pregnancy, lactation or women of childbearing age, weigh potential benefit against possible hazard.

Precautions: If combined with other psychotropics or anticonvulsants, consider carefully pharmacology of agents employed. Usual precautions indicated in patients severely depressed, or with latent depression, or with suicidal tendencies. Observe usual precautions in impaired renal or hepatic function. Limit dosage to smallest effective amount in elderly and debilitated to preclude ataxia or oversedation.

Side Effects: Drowsiness, confusion, diplopia, hypotension, changes in libido, nausea, fatigue, depression, dysarthria, jaundice, skin rash, ataxia, constipation, headache, incontinence, changes in salivation, slurred speech, tremor, vertigo, urinary retention, blurred vision. Paradoxical reactions such as acute hyperexcited states, anxiety, hallucinations, increased muscle spasticity, insomnia, rage, sleep disturbances, stimulation, have been reported; should these occur, discontinue drug. Isolated reports of neutropenia, jaundice, periodic blood counts and liver function tests advisable during long-term therapy.

Roche
ROCHE LABORATORIES
Division of Ciba-Geigy, Ltd., Nutley, N.J.
Roche, New Jersey 07110

www.decodog.com

History

- Widely available starting in the 1960s
- 1973 – 87 million prescriptions
- 1980s – high potency benzodiazepines found to be more effective for panic and anxiety than other drugs
 - Advantages: rapid onset and less risk of dependence
- 2007 – 74 millions prescriptions

Widespread Use

- Due to their significant margin of safety and effectiveness
 - BZDs are among the most prescribed psychotropic medications worldwide
 - On WHO essential drug list that should be available in all countries



DEA NFLIS 2006 Report


- Prescription drugs seized by law enforcement and analyzed forensics labs: 2001-2005

Drug	Rx Dispensed	Items seized per 10k Rx Dispensed
Diazepam	65M	6.06
Alprazolam	169M	5.96
Morphine	23M	5.80
Oxycodone	161M	5.29
Clonazepam	82M	3.55
Hydrocodone	550M	1.63
Codeine	165M	1.06

DEA NFLIS 2006 Report

- Prescription drugs seized by law enforcement and analyzed forensics labs

Figure 1. Distribution of benzodiazepines within region, 2001–2005.



Benzodiazepine Effects





Why benzos?

- Indications for a prescription:
 - Anxiety, insomnia, nausea, seizure, agitation, procedural sedation
- Advantages:
 - Immediate anxiolysis vs. buspirone and SSRI
- Other reasons:
 - Started for acute indication, yet not discontinued
 - Boosting other sedating medications (opioids)
 - Euphoria seeking
 - Opioid or alcohol withdrawal or cocaine toxicity

Mechanism of Action

- Organic bases with BENZene ring + DIAZEPENE moiety
- Modulates the gamma-aminobutyric acid A (GABA-A) receptor boosting affinity of GABA for the receptor
 - GABA - chief inhibitory neurotransmitter
- BZDs slow the brain down
- GABA receptor density low in respiratory brainstem > limiting the incidence of respiratory depression



Classes of Benzos

- Side chains determine potency, duration of action, and elimination
- Short-acting – Oxazepam
 - few active metabolites, clearance unaffected by age or liver disease
- Intermediate-acting – Lorazepam
- Long-acting – Diazepam and chlordiazepoxide
 - active metabolites, tissue accumulation, impaired clearance with age and liver disease

Tolerance

- To sedative and euphoric effects in days
- To anti-epileptic effects limits use for chronic seizure control
- Incomplete tolerance to cognitive impairment
- To the anxiolytic effects “is practically nonexistent”

Up To Date. Sedatives and hypnotics: Pharmacology and epidemiology
Principles of Addiction Medicine, 4th edition. P.105-6.

Adverse Effects


- Sedation
- Lethargy
- Respiratory Depression
- Impaired motor skills
- Impaired judgment
- Cognitive dysfunction
- Delirium
- Short-term memory impairment
- Anterograde amnesia
- Ataxia
- Hypotonia
- Depressed mood
- Exacerbation of COPD, sleep apnea

Patients often do not recognize their own impairment


BZDs in elderly

- Among Medicare enrollees, hip fracture linked to BZDs, regardless of half-life
 - First 2wks is highest risk time
 - Wagner et al. Arch Intern Med 2004;164; 1567.
- Beers Criteria 2002 for potentially inappropriate medication use in elderly
 - Medium/high doses of short acting BZDs and all long-acting BZDs
 - Fick et al. Arch Intern Med 2003; 163; 2716.

Does the cognitive function of long-term benzo users improve following withdrawal?



Are they still impaired compared to controls or normative data?




Barker. Arch Clinical Neuropsych 19 (2004) 437–454

Safety - Overdose

- Without concomitant sedating medications, benzos rarely cause life-threatening overdose
 - Sedated with normal vital signs
- Flumazenil – competitive antagonist of the GABA-A receptor that can reverse BZD actions
 - Risks of reversal (seizures and agitation) usually outweigh the benefits outside of procedural sedation


Principles of Addiction Medicine, 4th edition, chapter 47.
Up To Date. **Benzodiazepine poisoning and withdrawal**

Poisoning hospitalizations 1999-2006




Coben et al. Am J Prev Med. 2010.

Overdose deaths in NYC 2006-2008



NYC Vital Signs. NYC DPMH. 2010

Opioid overdose deaths in WVa 2006



Paulozzi et al. Addiction 2009; 104:1541.

Overdose and oversedation

Among 250 opioid dependent subjects with previous methadone or buprenorphine prescriptions, reported the following symptoms when also taking benzos:

Symptom	Methadone	BPN
Extreme drowsiness	42%	24%
Unconsciousness	7%	3%
Overdose	7%	1%

Nielsen. Addiction 2007; 102: 616-622.

Withdrawal Syndrome

- Symptoms? > Similar to alcohol withdrawal
 - Tremors, anxiety, perceptual disturbances, dysphoria, psychosis, seizures
- Onset of symptoms?
 - Varies by how long BZD has been used and half-life
 - can occur w/in hours for short-acting or weeks for long-acting

Treatment

Should I prescribe
benzodiazepines in the first
place?

Prescribers are ambivalent

On the one hand

- Rarely the abuse drug of choice
- Given the amounts prescribed, benzo abuse is “remarkably low”
- Benzos work fast with few side effects
- Benefit maintained over time

Schenck CH; Mahowald MW Am J Med 1996 Mar;100(3):333-7.

On the other hand

- Non-medical use very common
- Concerning subgroups
 - Other sedating meds
 - Elderly
 - Other addictions
- Hard to discontinue
- Does not improve long-term course of PTSD
- Co-morbid depression may worsen

Stevens, Pollack. J Clin Psychiatry 2005; 66s2: 21-27

Which hand?

Consider when prescribing benzos

- Intent –
 - Are you treating a diagnosed medical problem?
- Effect –
 - Does the medication improve the patient's functional status or worsen it?
- Monitoring –
 - Are you assessing the patient at the peak or trough effect of the medication?

Treatment

How do I get a patient off benzos,
yet minimize withdrawal and
relapse?

Withdrawing benzodiazepines

Don't stop BZDs abruptly due to risk of withdrawal seizures

Strategy 1 - Taper over weeks to months

- Taper 10% starting dose every 1-2 wks
- Decrease taper amount and lengthen interval for final 25-35% of taper.

Consider cognitive behavioral therapy/SSRI during and after for breakthrough symptoms

Principles of Addiction Medicine. 4th edition. P. 581-4.

Withdrawing benzodiazepines

Don't stop BZDs abruptly due to risk of withdrawal seizures

Strategy 2 - Substitute and taper

1. Stabilize on phenobarbital using hx and symptoms
2. Dose TID
3. Taper 30mg per day or slower

Drug	Phenobarbital 30mg equiv.
Alprazolam	0.5-1mg
Clonazepam	1-2mg
Diazepam	10mg
Lorazepam	2mg

Consider cognitive behavioral therapy/SSRI during and after for breakthrough symptoms

Principles of Addiction Medicine. 4th edition. P. 581-4, 649.

Withdrawing benzodiazepines

Don't stop BZDs abruptly due to risk of withdrawal seizures

Strategy 3 – Taper with adjunct

- Short BZD taper (3 days?) and treat with anti-convulsant for 2-3 weeks
 - carbamazepine 200 TID or
 - valproic acid 250TID

Consider cognitive behavioral therapy/SSRI during and after for breakthrough symptoms

Principles of Addiction Medicine. 4th edition. P. 581-4.

Pages and Ries. Am J Addictions 1998;7;198.

Harris et al. Alcohol Alcohol 2000;35; 319-325.

BZD treatment – Cochrane Review

Cochrane review authors conclusions:

1. Gradual taper is preferable to abrupt discontinuation
2. Carbamazepine appears to be a helpful adjunct to gradual taper
3. Confirming the benefit of switching from short to long-acting at beginning of taper requires further study

Denis et al. Pharmacological interventions for benzodiazepine **mono-dependence** management in outpatient settings. *Cochrane Database of Systemic Reviews*. 2006.

Case

- 29 yo man presents for follow-up at methadone clinic after inpatient admission for femur fracture from falling onto subway track from platform.
- Treated with clonazepam since age 16 for panic disorder. Takes 6mg in divided doses daily. Missed his afternoon dose, the day of the accident
- Started using heroin at age 23
- On methadone maintenance for 1 year, doing well, about to get his first take home

Thoughts

- Did BZDs cause his fall?
 - or did not taking BZDs cause his fall?
- Is he addicted to benzos?
- Should he come off of BZDs? How?
- If not, how do we keep him safe?
- Should he get take homes?
- Did teenage BZD treatment cause his heroin addiction?

What should be done about benzos?

- Prescribe with caution
- Educate patients
 - Safety first – Teens, mixing meds
 - Function over feelings
 - Risk of tolerance to benefits and withdrawal
 - Communication among prescribers
- Discontinue if risks outweigh the benefits

Thanks!

Alex Walley, MD, MSc
awalley@bu.edu


Benzos in Methadone Patients

Upon MMT entry in Israel


- 47% of patients abusing benzos ceased after 1 year
- 27% of patients not abusing benzos had started by 1 year
- Reasons for abuse included:
 - 87% to improve emotional state
 - 41% to boost other drugs
 - 40% for sleep
 - 24% to get high on benzos alone
 - 23% for withdrawal
 - 19% to reduce the effects of stimulants

Gelkopf et al. DAD 1999; 55: 63-68.


Acute to chronic opioid use



Methadone Maintenance



Boosting methadone with benzodiazepines



Patient's perspective on chronic benzo use

- Purpose of benzos
 - Means of coping with stress/anxiety and insomnia

“Sometimes my life gets so up in the air that I say to my children, right now I wish I had a wafer-sized Valium.”
 - Lifeline or life-transforming properties

“It makes me want to go on living.”
“I think if it weren't for the chemicals I wouldn't be chugging along.”
“I don't mentally think I would have survived without it and that's the truth.”
 - Lack of awareness, underestimation, disregard for side effects

“He wouldn't have given it to me if he thought it was gonna hurt me.”
“It's just a small, little, tiny white pill.”
“It's the lowest dose that they make.”
“My head always feels foggy.”

Cook et al. JGIM 2007: 22; 1094-1100.

Patient's perspective on chronic benzo use

- Attitudes toward taper/ discontinuation
 - Resistance to taper

“I see no reason why I should put myself through hell... We don't have that long to live and we might as well enjoy ourselves while we're here.”

“On numerous occasions I've tried to go off of it. And the reaction is I can't sleep and I'm totally wired. I'm up all night.”
 - Rejection of psychological interventions

“I just don't want to. I'm not one of those people who can sit around and talk about my problems with strangers.”
 - The physician-patient relationship

“I have complete faith in Dr. _____. I mean we go back a lot of years. Whatever he says, goes.”

Cook et al. JGIM 2007; 22; 1094-1100.

Benzos for panic disorder

Anxiolysis maintained – dose decreased




Fig 1.—Frequency distribution of daily alprazolam dose at discharge (open bars) and follow-up (FU) (closed bars) illustrates reduction in dose (36 [60%] of patients) or discontinuation of alprazolam (18 [30%] of patients) at FU. Mean \pm SEM dose was 3.1 ± 0.2 mg/d at time of discharge for total group and 1.2 ± 0.2 mg/d at FU (1.8 ± 0.2 mg/d for patients continuing to receive alprazolam at FU, discharge vs FU, both groups, $P < .001$, paired t test, two-tailed).




Fig 2.—Frequency of panic attacks at admission (ADM), discharge (DC), and follow-up (FU) for subgroups receiving (open bars) and not receiving (closed bars) alprazolam at FU. 1 indicates patients not receiving alprazolam at FU: ADM vs DC, $P < .05$, and ADM vs FU, $P = .01$, paired t test, two-tailed; 2, patients receiving alprazolam at FU: ADM vs DC, $P < .001$, and ADM vs FU, $P = .005$, paired t test, two-tailed; and 3, patients not receiving alprazolam at FU vs patients receiving alprazolam at FU: status at ADM, $P < .05$, and status at FU, $P < .05$, Student's t test, two-tailed.

Nagy et al. Arch Gen Psychiatry 1989; 46; 993.

N=60 – Natural history observation cohort

Benzos in Methadone Patients

Among 361 cocaine/heroin users enrolled in a contingency management trial, benzodiazepine use was associated with

- Increased cocaine use during treatment
- Blunted response to CM

Ghitza et al. DAD 2008; 97:150-157.

Prescription Drug Misuse

- Higher doses than prescribed
- More frequently than prescribed
- Without a prescription
- Reasons other than intended by the prescriber

Self-medication

- One physician survey reported that:
 - 26% of psychiatrists
 - 11% of other physicians
- Used unsupervised benzodiazepines in the past year

Principles of Addiction Medicine, 4th edition p.535.

Pharmacology & Kinetics


- Acts on GABA receptors by potentiating its inhibitory effects on the CNS
- Kinetics divided into three groups:
 - Short acting
 - Triazolam (Halcion), Oxazepam (Serax), Alprazolam (Xanax)
 - Midazolam (Versed) → but has more active metabolites
 - Intermediate acting
 - Lorazepam (Ativan), Temazepam (Restoril)
 - Long acting
 - Diazepam (Valium), Chlordiazepoxide (Librium), Clonazepam (Klonopin)
- Rapidly absorbed in GI tract, metabolized in liver

Withdrawal syndrome

Marked decrease or cessation of benzodiazepines after several weeks of regular use and 2 or more ...


- Autonomic hyperreactivity
- Increased hand tremor
- Insomnia
- Nausea or vomiting
- Hallucinations
- Psychomotor agitation
- Anxiety
- Tonic-clonic seizures

Poisoning hospitalizations 1999-2006



Coben et al. Am J Prev Med. 2010.


New Users - 2008 NSDUH




Nonmedical use – “not prescribed for the respondent or that the respondent took only for the experience or feeling that the drug caused”

Mean Age at First Use 2008

NSDUH



Past Month Use: 2002-2008



NSDUH 2008

Z drugs

- Zolpidem (ambien), esopiclone (lunesta) and zaleplon (sonata)
- Same GABA target
- Case reports of withdrawal and abuse
- Post-marketing surveillance indicates low abuse potential considering the amount prescribed

Urine testing

- Urine specimens contain little parent BZD
- Many immunoassays detect oxazepam: less likely clonazepam, lorazepam or triazolam unless present in high doses
- Chlorazepate, chlordiazepoxide, diazepam, and temazepam are metabolized to oxazepam

Metabolism

- Primarily hepatic, typically by CYP2C19 and/or CYP3A4
- Oxazepam, temazepam, and lorazepam are not metabolized by the liver, and excreted by the kidney

Up To Date. **Benzodiazepine poisoning and withdrawal**